

Analysis of a XMM-Newton Spectrum of the Extremely Hot White Dwarf in Nova V4743 Sgr

T. Rauch^{*†}, K. Werner[†] and M. Orio^{**‡}

**Dr.-Remeis-Sternwarte, Bamberg, University of Erlangen-Nürnberg, Germany*

† Institute for Astronomy and Astrophysics, University of Tübingen, 72076 Tübingen, Germany

***Istituto Nazionale di Astrofisica (INAF), Osservatorio Astronomico di Torino, Italy*

‡ Department of Astronomy, University of Wisconsin, Madison, USA

Abstract. Classical novae occur in close binary systems (main-sequence star + white dwarf). V4743 Sgr is a very fast nova exhibiting a X-ray spectrum which is consistent with a view on the optically thick atmosphere of an extremely hot white dwarf (WD). It exhibits strong absorption features of the C V, C VI, N VI, N VII, and O VII resonance lines indicating an effective temperature of about 610 kK.

We present preliminary results of a spectral analysis by means of line-blanketed non-LTE atmosphere models of a XMM-Newton RGS-1 spectrum of V4743 Sgr, taken about half a year after its outburst.

V4743 Sgr has been discovered as a nova on Sep 20, 2002 (Haseda et al. 2002). Until Mar 2003, it evolved into the brightest super-soft X-ray source in the sky. A first CHANDRA spectrum shows a flux maximum around 30 Å and strong C V, N VI, and N VII (Fig. 1) absorption features which suggest temperatures of 1 – 2 MK (Ness et al. 2003). These lines are blue shifted (-2400 km/sec) – most likely we see expanding gas which consists out of CNO-processed material.

However, in the phase after the outburst, i.e. when the H burning is still on-going (for years), the surface composition of the WD is poorly known. In order to make some progress, we decided to use our static NLTE models to investigate basic parameters like effective temperature (T_{eff}) and surface composition. Although we neglect any effect of the velocity field, our aim is at least a qualitative modeling of the XMM RGS-1 spectrum of V4743 Sgr, obtained on Apr 4, 2003 with an exposure time of about 10 h. The EPIC-pn count rate in timing mode was 1348 cts/sec (Orio et al. 2003).

For a preliminary analysis, we employed plane-parallel, static models which are calculated with TMAP, the Tübingen NLTE Model Atmosphere Package (Werner et al. 2003, Rauch & Deetjen 2003). The first set of models is composed of H+He+C+N+O, then Ne+Mg+S are added subsequently in a line-formation calculation (Tab. 1). However, in general TMAP can treat all elements from hydrogen to the iron group simultaneously (Rauch 1997, 2003).

Small grids of model atmospheres were calculated in order to match the relevant T_{eff} range (0.5 – 1 MK). 208 levels of H+He+C+N+O+Ne+Mg+S are treated in NLTE with 332 individual line transitions (Tab. 1).

A comparison with the spectrum of V4743 Sgr via XSPEC yields a T_{eff} of about 705 kK, however, the flux at wavelengths smaller than 26 Å is too low. Decreasing the C

TABLE 1. Statistics of the model atoms used in our calculations of the NLTE model atmospheres. The notation is: NLTE = levels treated in NLTE, LTE = LTE levels, RBB = radiative bound-bound transitions

atom	ion	NLTE	LTE	RBB	atom	ion	NLTE	LTE	RBB
H	I	5	11	10	Ne	VII	10	50	12
	II	1	-	-		VIII	8	18	15
He	I	1	21	0	Mg	IX	5	6	3
	II	10	22	45		X	1	0	0
	III	1	-	-		VIII	1	19	0
C	V	29	21	60		IX	3	23	1
	VI	15	21	26		X	2	3	1
	VII	1	0	-		XI	5	6	1
N	V	5	15	6	S	XII	1	0	0
	VI	17	7	33		X	1	7	0
	VII	15	21	30		XI	5	29	1
	VIII	1	0	-		XII	10	12	15
O	VI	1	34	0		XIII	9	21	10
	VII	19	7	21		XIV	9	1	15
	VIII	15	30	30		XV	1	0	0
	IX	1	0	-					
total							208	409	332

abundance by a factor of 0.01, results in a much better fit in this region (Fig. 1).

A crucial point for the understanding of processes during the nova outburst may be the time-dependent prediction of surface abundances as well as the spectral analysis of high-resolution X-ray spectra taken from outburst to the end of surface H burning. The inspection of available XMM-Newton spectra of V4743 Sgr, taken on a half-year time scale has shown that about weekly spectra are highly desirable. This would also allow to investigate the velocity reliably.

Starrfield et al. (2000) calculated the yields of thermonuclear runaways of O/Ne/Mg WD. We adopted the average of their models' abundance ratios and calculated a grid of test models. Their fluxes do not reproduce the observation better than our simple C-reduced models. Thus, this issue will be a challenge for both, theory and spectral analysis in the future.

The comparison of our preliminary model atmosphere fluxes with the Apr 2003 XMM-Newton RGS spectrum of V4743 Sgr, however, has shown that at solar abundance ratios, we cannot fit the overall flux distribution. Although the fit is not perfect yet, it is obvious that the C/N abundance ratio is much smaller (about a hundred times) than the solar value, typical for CNO processed material. The fit to the observation based on our first test models is best at a of T_{eff} about 610 kK and $\log N_{\text{H}} = 20.54 \text{ [cm}^{-2}\text{]}$.

Analysis of X-ray data in Tübingen is supported by the DLR under grant 50 OR 0201.




FIGURE 1. Comparison of the XMM-Newton RGS-1 spectrum with NLTE model fluxes (thick: H:He:C:N:O = 18.9:75.1:0.0006:0.016:5.96, thin: solar). T_{eff} and N_{H} are determined with XSPEC. A black body (dashed) is shown for comparison only and to identify instrumental features of XMM-Newton. The synthetic spectra are shifted by -2400 km/sec

REFERENCES

1. Haseda, K., West, D., Yamaoka, H., Masi, G. 2002, IAU Circ. 7975
2. Kato, T., Hishikura, T., West, J.D., et al. 2002, IAU Circ. 7976
3. Ness, J.-U., Starrfield, S., Burwitz, V., et al. 2003, ApJ 594, L127
4. Orio, M., Leibowitz, E., Rodriguez, P., et al. 2003, IAU Circ. 8131
5. Rauch, T. 1997, A&A 320, 237
6. Rauch, T. 2003, A&A 403, 709
7. Rauch, T., Deetjen, J.L. 2003, in: Workshop on Stellar Atmosphere Modeling, eds. I. Hubeny, D. Mihalas, K. Werner, The ASP Conference Series (San Francisco: ASP), 288, 103
8. Starrfield, S., Sparks, W.M., Truran J.W., Wiescher, M.C. 2000, ApJS 127, 485
9. Werner, K., Dreizler, S., Deetjen, J.L., Nagel, T., Rauch, T., Schuh, S.L. 2003, in: Workshop on Stellar Atmosphere Modeling, eds. I. Hubeny, D. Mihalas, K. Werner, The ASP Conference Series (San Francisco: ASP), 288, 31